

# Iodine Status in Pregnant Women in the National Children's Study and in U.S. Women (15–44 Years), National Health and Nutrition Examination Survey 2005–2010

Kathleen L. Caldwell,<sup>1</sup> Yi Pan,<sup>1</sup> Mary E. Mortensen,<sup>1</sup> Amir Makhmudov,<sup>1</sup> Lori Merrill,<sup>2</sup> and John Moyer<sup>3</sup>

**Background:** This report presents iodine data from National Health and Nutrition Examination Survey (NHANES) and from a sample of pregnant women in the National Children's Study (NCS) Vanguard Study.

**Methods:** Urinary iodine (UI) was measured in a one third subsample of NHANES 2005–2006 and 2009–2010 participants and in all 2007–2008 participants age 6 years and older. These measurements are representative of the general U.S. population. UI was also measured in a convenience sample of 501 pregnant women enrolled in the NCS initial Vanguard Study from seven study sites across the United States.

**Results:** NHANES median UI concentration in 2009–2010 (144  $\mu\text{g/L}$ ) was significantly lower than in 2007–2008 (164  $\mu\text{g/L}$ ). Non-Hispanic blacks had the lowest UI concentrations (131  $\mu\text{g/L}$ ) compared with non-Hispanic whites or Hispanics (147 and 148  $\mu\text{g/L}$ , respectively). The median for all pregnant women in NHANES 2005–2010 was less than adequate (129  $\mu\text{g/L}$ ), while third trimester women had UI concentrations that were adequate (median UI 172  $\mu\text{g/L}$ ). Third trimester women participating in the NCS similarly had an adequate level of iodine intake, with a median UI concentration of 167  $\mu\text{g/L}$ . Furthermore, NCS median UI concentrations varied by geographic location.

**Conclusions:** Dairy, but not salt, seafood, or grain consumption, was significantly positively associated with median UI concentration in women of childbearing age. Pregnant women in their third trimester in the NHANES 2005–2010 had adequate median UI concentrations, but pregnant women in NHANES who were in their first or second trimesters had median UI concentrations that were less than adequate. Non-Hispanic black pregnant women from both the NHANES 2005–2010 and the NCS consistently had lower UI median concentrations than non-Hispanic whites or Hispanics.

## Introduction

ADEQUATE DIETARY IODINE INTAKE is essential to the synthesis of thyroid hormone, which is key for normal growth, development, and metabolism throughout life. Recent data show that 1.88 billion people globally, including 241 million school children, have insufficient dietary iodine intake (1). Iodine deficiency disorders include a range of health consequences or abnormalities of the body resulting from a prolonged lack or insufficient intake of iodine, ranging from simple goiter to cretinism, which is a condition of severely stunted physical and mental growth (1,2). Adequate iodine intake during pregnancy and lactation is especially critical to normal brain development in the fetus (3). While

diet-induced hypothyroidism can occur at any stage of life, the most devastating consequences of iodine deficiency occur during fetal development and early childhood, and include miscarriage, stillbirth, congenital abnormalities, and severe and irreversible mental retardation. Iodine deficiency is the leading and most preventable cause of mental retardation in the world (4). The consequence of severe iodine deficiency is a 10%–15% reduction in IQ for a population (4).

Strategies to combat the prevalence of iodine deficiency disorders focus on ensuring adequate dietary intake. Dairy, grain, seafood, and to a lesser degree, iodized salt are the major sources of iodine in the United States (5). However, there is a wide variation in iodine content in foods and iodine content is rarely included in nutritional labeling (5). In the

<sup>1</sup>Division of Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia.

<sup>2</sup>Westat, Rockville, Maryland.

<sup>3</sup>National Children's Study Program Office, Eunice Kennedy Shriver National Institute for Child Health and Human Development, National Institutes for Health, Bethesda, Maryland.

United States approximately 60% of iodine consumed comes from dairy products. The iodine is added to dairy products as a consequence of iodine added to cattle feed (6), or use of iodophor disinfectants in the milking process (7,8). Iodized salt accounts for a small amount of dietary iodine. Approximately 70% of salt consumed in the United States comes from processed and restaurant foods which generally do not use iodized salt (9).

The Institute of Medicine has set the recommended dietary allowance (RDA) for iodine in adult men and women at 150  $\mu\text{g}$  per day (10). One teaspoon of iodized table salt contains approximately 400  $\mu\text{g}$  of iodine. To support fetal and infant thyroid function, the Institute of Medicine suggests an RDA for pregnant women of 220  $\mu\text{g}$  iodine per day and 290  $\mu\text{g}$  iodine per day during breastfeeding (10). To prevent iodine deficiency, the American Thyroid Association recommends supplementation containing 150  $\mu\text{g}$  of iodine daily for U.S. women of childbearing age during the preconception phase, as well as during pregnancy and lactation (11). In addition, adequate iodine intake before conception (RDA of 150  $\mu\text{g}$  per day) is important to ensure adequate maternal iodine stores to support the fetus. Preconception iodine status of the mother influences the degree of successful maturation of the fetal central nervous system and subsequent neurodevelopment of the child (12).

There are no analytical techniques available to directly measure an individual's daily iodine status. Urinary iodine (UI) reflects iodine intake within the past few days. As a clinical biomarker, UI generally is not useful to classify intake sufficiency or deficiency in a person, but rather to define the risk of a population. UI measurement in 24-hour urine collection is preferred, but iodine excretion can be expressed per gram of creatinine in spot urine collections in population groups with very low inter- and intra-individual variation in urinary creatinine (13). The World Health Organization (WHO) defines nutritional iodine sufficiency for a population by UI concentrations as follows: excessive iodine intake,  $>300 \mu\text{g}/\text{L}$ ; more than adequate intake,  $200\text{--}299 \mu\text{g}/\text{L}$ ; adequate intake,  $100\text{--}199 \mu\text{g}/\text{L}$ ; mild iodine deficiency,  $50\text{--}99 \mu\text{g}/\text{L}$ ; moderate iodine deficiency,  $20\text{--}49 \mu\text{g}/\text{L}$ ; and severe iodine deficiency  $<20 \mu\text{g}/\text{L}$ . The WHO recently defined adequate iodine intake for pregnant women as an iodine excretion of  $150\text{--}249 \mu\text{g}/\text{L}$  and inadequate iodine intake as represented by iodine excretion  $<150 \mu\text{g}/\text{L}$  (13).

Given the significant numbers at risk for iodine insufficiency among select population subgroups in the United States, population-wide monitoring to document UI levels is a necessary public health activity. Recent U.S. population data from the National Health and Nutrition Examination Survey (NHANES) have shown that women and some subpopulations of non-Hispanic blacks tend to have UI concentrations indicating mild iodine deficiency (14). The National Children's Study (NCS) initial Vanguard Study was launched during the same time frame as the 2009–2010 NHANES. The NCS Vanguard Study was intended to provide operational data to inform decisions regarding processes and measures for inclusion in the Main Study. The Centers for Disease Control and Prevention (CDC) used samples collected during the Vanguard Study to conduct a pilot study. The purpose of the CDC pilot study was to measure a number of environmental chemicals and nutritional biomarkers in the NCS convenience sample of pregnant women from seven study


locations throughout the United States. As part of the CDC pilot study, UI was measured in Vanguard Study participants who were in their third trimester of pregnancy. These data help provide a more complete picture of UI status of U.S. pregnant women and enhance the data provided by the NHANES. Our paper will focus on iodine status in the U.S. population as measured by the most recent NHANES 2009–2010 cycle and pregnant women from NHANES 2005–2010 as well as the UI levels for pregnant women who participated in the NCS Vanguard pilot study.

UI has been measured in NHANES since 1971. In recent years, UI has shown a decrease. In all instances in which the data are stratified by race/ethnicity, the non-Hispanic black population has had the lowest UI relative to other racial/ethnic groups. Despite oversampling of pregnant women in NHANES 2001–2006, the number of pregnant women participating in the survey was still quite small. The CDC collaboration with the NCS provided an opportunity to measure UI in a large number of third trimester pregnant women as part of a pilot study to evaluate the clinical biospecimen collection protocols used in the 2009–2010 Vanguard Study. The Division of Laboratory Sciences, National Center for Environmental Health measured UI concentrations and other environmental and nutritional analytes in samples collected from participants enrolled at any of the seven original Vanguard study sites. In this study, results of UI measured in the NCS sample of pregnant women provides additional data on a group at risk for inadequate iodine intake and will be compared with results from a representative sample of U.S. pregnant women from NHANES.

## Data and Methods

### *NHANES design*

NHANES is conducted by the National Center for Health Statistics (NCHS) of the CDC. It is a program of studies designed to assess the health and nutritional status of adults and children in the United States. The NHANES interview includes demographic, socioeconomic, dietary, and health-related questions. Physical examinations and sample collection for laboratory tests are included. The survey is conducted in 2-year cycles. In this article, the NHANES 2009–2010 survey cycle provides the main source of population data for UI. However, we utilized dietary data from NHANES 2007–2008 (dietary data for 2009–2010 were not yet available) and included UI results for pregnant women in NHANES 2005–2010 in order to have a larger sample size. The survey incorporates sample population weights to account for the unequal selection probabilities caused by the cluster design, nonresponse, and planned oversampling of certain subgroups. Pregnant women were oversampled in NHANES 2001–2006. Due to the relatively small number of pregnant women in each 2-year cycle for some analyses, we combined UI results for pregnant women from NHANES 2005–2010 for different analyses in this report. For pregnant women ages 15–44 years, UI results from NHANES 2007–2010 were used. For pregnant women ages 15–44 years by trimester, UI results from NHANES 2005–2010 were used. For the entire population, UI results from NHANES 2009–2010 were used. Figure 1 illustrates the combinations of data across several cycles that were used for the projects and a justification for use of different cycles is given.



**FIG. 1.** Description of National Health and Nutrition Examination Survey (NHANES) and National Children’s Study (NCS) population utilized for each of the different analyses in this investigation and the justification.

Data on the sociodemographic variables sex, age, race/ethnicity, and pregnancy status were collected. Age was categorized using the following groups: 6–11, 12–19, 20–29, 30–39, 40–49, 50–59, 60–69, and 70 years and older. Race/ethnicity categories were self-reported as non-Hispanic white, non-Hispanic black, and Hispanic, which includes Mexican American and other Hispanic. Childbearing age was defined as between 15 and 44 years old. Pregnancy status was determined by a urine pregnancy test. Dairy consumption in the past 30 days was categorized as “never or rare,” “not often,” or “often.” Never or rare consumption meant any dairy product consumed less than once a week in the past 30 days; not often denoted some dairy product consumed more than once a week but less than once a day; and often meant dairy product was consumed more than once a day. Salt con-

sumption was determined by response to the question “How often is ordinary salt or seasoned salt added in cooking or preparing foods in your household? Is it never or rarely, occasionally or very often?” Fish or shellfish consumption during the past 30 days was also included for NHANES 2007–2008. Fish included breaded fish, tuna, bass, catfish, flatfish, haddock, mackerel, perch, pike, pollock, porgy, salmon, sardines, sea bass, swordfish, trout, and other fish. Shellfish consisted of clams, crabs, crayfish, lobsters, mussels, oyster, scallops, shrimp, and other shellfish. Total grain intake was determined based on intake reported on the first day of the dietary interview. Dietary supplement use during the past 30 days was determined by response to the question, “Have you used or taken any vitamins, minerals or other dietary supplements in the past month?”

### NCS design

NCS is a planned large long-term study of children's health and development. It examines the effects of the environment, as broadly defined to include factors such as air, water, diet, sound, family dynamics, community and cultural influences, and genetics on the growth, development, and health of children across the United States, following them from before birth until age 21 years. The goal of the study is to improve the health and well-being of children and contribute to understanding the role of various factors on health and disease. Details of the background and organization of this complex undertaking have been described (15,16). Briefly, the NCS structure consists of the Vanguard Study, Main Study, and formative research that supports and informs the Main Study. The Vanguard Study is a parallel extensive feasibility study that will precede the Main Study to evaluate feasibility, acceptability, and cost of recruitment, logistics, operations, and study visits. The original Vanguard Study began in 2009 and included recruitment of pregnant women at seven selected study locations throughout the United States. Women were eligible if they were not surgically sterile and were age 18 through 49 years or currently pregnant. The women had up to two pregnancy visits, which included interviews and collection of biological and environmental samples. Pregnancy visits were scheduled to occur in the first and third trimesters; this study used samples collected from women in their third trimester of pregnancy. Race/ethnicity categories were self-reported as non-Hispanic white, non-Hispanic black, and all Hispanic, which included Mexican American and other Hispanic. At the time of this submission, such interview survey results as dietary recall, food intake, and supplement use were not available.

### Sample collection

During the NHANES physical examinations, spot urine specimens were collected into a prescreened collection cup from participants ages 6 years and older, and aliquots of these specimens were generated and stored cold (2°C to 4°C) or frozen until shipped. The samples were shipped on dry ice to the CDC National Center for Environmental Health (NCEH) and were stored frozen (-70°C) for <1 year. During the NCS third trimester visits, spot urine specimens were collected from participants into a prescreened collection cup, frozen, and shipped to the NCS Repository (Fisher Bioservices, Rockville, MD) on dry ice, thawed, and aliquoted into prescreened metal-free cryovials, according to NCS protocols, then stored frozen at vapor phase liquid nitrogen temperatures until shipment on dry ice to the CDC NCEH laboratory where they were also stored frozen (-70°C) for <1 year.

### Iodine measurements

In all NHANES cycles except 2007–2008, UI measurements were obtained for one third of the survey population. For NHANES 2007–2008, UI measurements were obtained for all participants over age 5 years. Samples were analyzed for UI concentration using the method of Caldwell *et al.* (17,18). Briefly, 0.5 mL of urine was diluted 1:10 with 1% (v/v) tetramethylammonium hydroxide, 0.02% Triton<sup>TM</sup> X-100 (Mallinckrodt Baker, Inc. Phillipsburg, NJ), 25 µg/L tellurium, 5 µg/L bismuth, 5% (v/v) ethanol, 1000 µg/L gold, and 0.5 g/L EDTA. This solution was subsequently analyzed using in-

ductively coupled plasma dynamic reaction cell mass spectrometry. Iodine was quantified based on the peak as a ratio of analyte to internal standard tellurium. The limit of detection was found to be 1.4 µg/L. The LOD as reported is equal to 3×S/N, where S/N is the signal to noise derived from the measurement process as the concentration approaches zero. Four concentration levels of quality controls were analyzed in each analytical batch. Reported results met the accuracy and precision specification of the quality control/quality assurance program of the Division of Laboratory Sciences, NCEH, CDC (19). Two quality control pools were analyzed from 2000 to 2012. One QC pool ( $n=869$ ) with a concentration of 93 µg/L had a relative standard deviation (RSD) of 2.3%. The RSDs for the second pool ( $n=866$ ) with a concentration of 308 µg/L was 3.0%. Absolute assay accuracy was verified by the blind analysis of two additional iodine reference solutions and the analysis of National Institute of Standard Technology (NIST) 2670A, 2672a, and 3668 Standard Reference Materials.

Stability was excellent with no trend observed for the RSD across time. To confirm agreement between survey cycles, 122 previously analyzed urine iodine samples from NHANES study years 2000–2009 were reanalyzed in 2011 by the CDC lab to assure the repeatability of those measurements across time. Concordance correlation coefficient (CCC) (20) was applied to measure the agreement between the measurements of the same iodine sample among different time periods. Overall, the CCC was 0.9955 (95% confidence interval [CI 0.9948, 0.9962]), which assures the validity of the measurements from instruments across time. The stability of measuring the iodine samples by the instruments across time from 2000 to 2010 at CDC was also established through successful administration and participation in CDC's external quality assurance program: Ensuring the Quality of Urinary Iodine Procedures (21).

### Creatinine measurements

NHANES and NCS urinary creatinine concentrations were determined using the Roche/Hitachi Modular P Chemistry Analyzer (Laboratory Services, LLC, Hartford, CT) in 2009–2010 (19). This method is described in Roche's Creatinine plus Product Application no. 11775685216V18. Iodine concentrations were adjusted by using creatinine concentrations to correct for variable urine excretion rates at the time of spot urine specimen collection.

### Statistical analysis

Statistical analyses were conducted using SAS, version 9.2 (SAS Institute, Inc., Cary, NC), and SUDANN PROC DEScript, version 10.0 (Research Triangle Institute, Research Triangle Park, NC). In NHANES, median UI measurements and the median creatinine corrected UI concentrations with CIs were analyzed based on the method of Korn and Graubard (22). In each NHANES survey period we used sample weights to account for differential nonresponse or no coverage and to adjust for oversampling of some groups. Median test was used to compare the median UI concentrations between or among different target subpopulations in the NHANES analyses (23). Error proportions associated with certain UI thresholds were estimated and were flagged when the relative standard error was greater than 30%. In NHANES, Rao–Scott *F*-adjusted chi-square test was used to

test the association between two categorical variables. In NCS, median UI concentrations were analyzed for subpopulations and the Kruskal–Wallis test was applied to test for equality among different study sites. Histogram plots of iodine distribution on pregnant women of childbearing age and pregnant non-Hispanic black women of childbearing age were presented with both NHANES and NCS datasets.

**Results**

*U.S. population in NHANES 2009–2010*

From NHANES 2009–2010, median UI for the U.S. population 6 years and older was 144 µg/L [CI 132–154] (Table 1), which was significantly lower ( $p=0.001$ ) than the median of 164 µg/L [CI 154–173] in NHANES 2007–2008 (14). The median UI for all females was 134 µg/L [CI 125–146]. Median UI for women of childbearing age (15–44 years) was 124 µg/L [CI 111–139]. Median UI for non-Hispanic blacks was 131 µg/L [CI 120–144]. Non-Hispanic blacks had significantly lower UI concentrations compared to non-Hispanic whites and all Hispanics ( $p=0.03$ ; Table 1). As shown in Figure 2, a U-shaped curve with higher median UI at lower and higher age categories was observed in NHANES 2009–2010. Lowest UI was observed in ages 12–49 years, highest in children ages 6–11 years. The UI concentrations were consistently lower in women of childbearing age compared with the general population, a relationship that has been observed since 2001 (14) (Fig. 3).

*NHANES 2007–2010 U.S. pregnancy data and NCS*

Table 2 presents the median UI with CI in NHANES 2007–2010 by pregnancy status for women of childbearing age (15–44 years) and for NCS pregnant women. The median UI for pregnant women in NHANES 2009–2010 was 135 µg/L [CI 108–172] compared with 167 µg/L [CI 151–185] in third trimester pregnant women in the NCS study.

*Trimester data NHANES and NCS*

Given the limited number of pregnant women in both NHANES 2007–2010 ( $n=76$ ) or NHANES 2009–2010 ( $n=22$ ), stratification by trimester was only possible by using all pregnant women who participated in 2005–2010 NHANES (the number of pregnant women was 206; however, only 176 participants had trimester data available). Table 3 presents the median UI concentrations for women of childbearing age and pregnant women. Women in their first trimester had a median UI of 109 µg/L ( $n=42$ ) while the second trimester women had a median UI of 128 µg/L ( $n=70$ ). The median UI for NHANES women in their third trimester was 172 µg/L [CI 109–267] ( $n=64$ ), comparable to the NCS third trimester pregnant women (167 µg/L [CI 151–185],  $n=501$ ; Table 2).


*Distribution of UI in NCS and NHANES (2007–2010)*

Among NHANES pregnant women, approximately 55.8% ( $\pm 7.7\%$ ) had UI that suggested less than adequate iodine intake ( $<150$  µg/L, Table 2). Among NCS women, approximately 45.3% ( $\pm 2.3\%$ ) had UI that indicated less than adequate iodine intake. The distributions of UI for pregnant women in NHANES 2007–2010 and in NCS are presented in Figures 4A and 4B. Figure 4A shows the median UI of 135 µg/L and distribution in pregnant women from NHANES 2007–2010. The 5th and 95th percentiles of the UI distribution were 40 and 508 µg/L, respectively (Fig. 4A). Figure 4C shows the distribution for NHANES 2007–2010 pregnant non-Hispanic black women. The median UI concentration was 119 µg/L with 5th and 95th percentiles of 34 and 347 µg/L, respectively. Figures 4B and 4D show the distribution of UI in NCS women. The median UI for all NCS pregnant women was 167 µg/L, with 5th and 95th percentiles of 43 and 605 µg/L (Fig. 4B). For non-Hispanic black NCS pregnant women, median UI was considerably lower (132 µg/L) relative to all NCS pregnant women (Fig. 4D). It is interesting to note that only the NCS

TABLE 1. URINARY IODINE AND IODINE CORRECTED FOR CREATININE LEVELS BY AGE, SEX, AND RACE/ETHNICITY; WEIGHTED SURVEY ANALYSIS, NHANES 2009–2010

	n	Weighted n	Median [CI]	
			UI (µg/L)	I/Cr (µg/g creatinine)
Age, years				
6 and above	2864	271,924,206	144 [132–154]	143 [136–148]
6–11	379	23,034,180	213 [188–234]	250 [221–279]
12–19	454	32,605,343	131 [112–151]	117 [107–130]
20–29	328	41,055,524	132 [109–152]	112 [98–129]
30–39	361	39,145,156	132 [114–156]	114 [102–129]
40–49	368	42,930,574	128 [108–146]	114 [101–137]
50–59	300	40,067,968	136 [122–160]	147 [128–170]
60–69	316	27,097,884	148 [124–171]	181 [154–210]
70 and above	358	25,987,578	182 [160–206]	239 [195–264]
Sex				
Male	1402	132,645,050	152 [139–171]	135 [126–142]
Female	1462	139,279,157	134 [125–146]	150 [143–162]
Race/ethnicity				
Non-Hispanic white	1224	176,420,426	147 [133–160]	149 [143–159]
Non-Hispanic black	545	32,383,034	131 [120–144]	98 [89–105]
All Hispanic	914	41,172,201	148 [134–160]	142 [137–149]
Women of childbearing age	605	61,583,168	124 [111–139]	117 [108–127]

UI, urinary iodine; I/Cr, iodine corrected for creatinine levels; CI, 95% confidence interval.



**FIG. 2.** Median urinary iodine (UI,  $\mu\text{g/L}$ ) by age category, NHANES 2009–2010, participants 6 years and older. The solid horizontal reference lines indicate the adequate iodine intake range defined by the World Health Organization (WHO; 100–199  $\mu\text{g/L}$ ) for the general population. The dashed horizontal reference lines indicate sufficient median iodine intake for pregnant women (150–249  $\mu\text{g/L}$ ).

population has women with UI concentration greater than  $\sim 900 \mu\text{g/L}$ . These apparent outliers represent only few participants who are taking iodine-containing supplements or another significant source of iodine.


*Race/ethnicity in 2009–2010 NHANES and NCS*

Table 4 provides details on UI concentration by race/ethnicity and age from both 2009–2010 NHANES and NCS. Non-Hispanic blacks in each age category and in both study cohorts had lower median UI concentrations than non-Hispanic whites and all Hispanics. The trend is consistent for uncorrected and creatinine-corrected data.

*Geographic location (in NCS data only)*

Table 4 also provides the NCS sample population stratified by geographic locations. The NCS participants consist of

pregnant women from seven different geographic locations in the United States. The participating study centers were located in California, North Carolina, Minnesota and South Dakota, New York, Pennsylvania, Utah, and Wisconsin. Table 3 provides NCS median UI concentrations by the geographic region of the study site. All seven sites had different median UI concentrations. All collection sites used collection material that had been prescreened at the CDC. It is unlikely that protocol or material influenced the results. Urine iodine is very stable once collected as discussed in the Data and Methods section. Several study sites had median UI concentrations  $< 150 \mu\text{g/L}$ . The California median UI was  $107 \mu\text{g/L}$  well below those found in North Carolina, South Dakota-Minnesota, and Utah (217, 205, and  $190 \mu\text{g/L}$ , respectively). New York and Wisconsin had median UI concentrations of  $150 \mu\text{g/L}$  and  $145 \mu\text{g/L}$ , respectively, while Pennsylvania had a median UI of  $125 \mu\text{g/L}$ . Median UI concentrations in



**FIG. 3.** Median UI ( $\mu\text{g/L}$ ) for participants older than 6 years and women of child-bearing age based on NHANES 2001–2010.

TABLE 2. MEDIAN AND LOW URINARY IODINE CONCENTRATIONS IN U.S. WOMEN OF CHILDBEARING AGE (15–44 YEARS) BY PREGNANCY STATUS, NHANES 2007–2010 COMPARED TO NCS

	n	Median [CI]	<150 (µg/L)				<100 (µg/L)				<50 (µg/L)	
			%	SE	n	%	SE	n	%	SE	n	
NHANES total	1568	133 [121–143]	56.3	2.2	862	36.9	1.9	550	15.4	1.8	205	
NHANES Pregnant	76	135 [109–172]	55.8	7.7	45	26.3	6.5	26	15.7	6.6 <sup>a</sup>	13	
NHANES Nonpregnant	1492	133 [121–143]	56.3	2.2	817	37.3	1.9	524	15.4	1.9	192	
NCS Pregnant	501	167 [151–185]	45.3	2.3	227	26.7	1.9	134	7.0	1.5	35	

<sup>a</sup>The relative standard error >30%. The NHANES guidelines recommend a relative standard error ≤30%. SE, standard error.

California, Pennsylvania, and Wisconsin were below the recommended median UI concentration of 150 µg/L for pregnant women.

*Dietary questions in NHANES 2007–2008*

Dairy products consumption has been shown to be significantly associated with iodine intake (6). Table 5 shows the association between dairy consumption in the past 30 days and race/ethnicity for women of childbearing age in NHANES 2007–2008. The proportion of non-Hispanic black women who never or rarely consumed dairy products in the past 30 days was 39.8%, significantly higher than non-Hispanic whites (27.1%) or all Hispanics (31.1%) (*p*=0.004, Rao–Scott *F*-adjusted chi-square test). The median UI concentrations for non-Hispanic blacks were the lowest in each dairy consumption level compared to other race/ethnicity groups.

The relationship between salt consumption and median UI was also evaluated across race/ethnicity categories (Table 5). Hispanics reported greater salt use than the other groups, with 17.2% reporting never or rare use during the last 30 days compared to 24.5% for non-Hispanic blacks and 27.1 % for non-Hispanic whites. The proportion of Hispanic women who reported using salt often was 55.6 %, significantly higher than non-Hispanic blacks (45.9%) and non-Hispanic whites (35.9%) (*p*<0.0001, Rao–Scott *F*-adjusted chi-square test).

Fish or shellfish intake during the last 30 days was not significantly associated with median UI concentrations in racial/ethnic groups with the exception of Hispanics (Table 5). Hispanic women who reported fish or shellfish intake had significantly higher median UI compared to non-fish-consuming Hispanics (*p*=0.03). There was no significant association between higher median UI and grain intake, or

supplement use in each race/ethnicity category during the past 30 days (data not shown).


**Discussion**

NHANES 2009–2010 indicates that median UI in the general population is 144 µg/L [CI 132–154]. That concentration is significantly lower than in the 2007–2008 cycle, in which median UI was 164.0 µg/L [CI 154–173] (*p*=0.001). In NHANES 2009–2010, certain groups within the U.S. population did not achieve adequate dietary iodine intake. Non-Hispanic blacks had a significantly lower UI concentration than non-Hispanic whites and all Hispanic groups. Non-Hispanic white children had the highest UI levels, in the more than adequate range, while non-Hispanic black and Hispanic children had median UI in the adequate range. In the U.S. population, age groups 12–49 years had the lowest UI, which has been noted before (14). As in previous NHANES survey periods, women had lower UI concentrations than men (14). NHANES data (2007–2010) indicated that 37.3% of nonpregnant women of childbearing age (*n*=1492) have UI concentrations below 100 µg/L, while 55.8% of pregnant women (*n*=76) have UI concentrations less than 150 µg/L. Meanwhile, in the convenience sample of 501 pregnant women from the NCS, 45.3% of the participants had UI concentrations less than 150 µg/L. The observed difference in the proportion of pregnant women with UI<150 µg/L is most likely because the NCS participants were all in their third trimester of pregnancy. While there were only 76 pregnant women in NHANES 2007–2010 with UI measurements, limiting our ability to look at data stratified by trimester of pregnancy due to insufficient power, by combining NHANES 2005–2010 data we were able to increase our sample of pregnant women to 206. We stratified these women into trimester groups. The NHANES third trimester UI concentration 172 µg/L [CI 109–267] is very similar to the UI concentration found in the third trimester women participating in the NCS pilot study (167 µg/L [CI 151–185]).

UI levels found in women in the third trimester in both NHANES and NCS were in the adequate iodine intake range. Women in late pregnancy may be more attuned to maintaining adequate nutrition or may be taking supplements with iodine. Most iodine-containing multivitamins should have at least 150 µg of iodine, but only about half of the multivitamins sold in the United States contain iodine (12). According to Gregory *et al.* (12), in the NHANES 2001–2006, approximately 76.9% of pregnant women reported prenatal supplement use. However, only 20.3% of those preparations

TABLE 3. MEDIAN URINARY IODINE CONCENTRATIONS FOR WOMEN OF CHILDBEARING AGE AND PREGNANT WOMEN BY TRIMESTER, NHANES 2005–2010

Category	n	UI (µg/L) median [95% CI]
Total	2233	129 [120–136]
Pregnant	206	129 [101–173]
Nonpregnant	2027	129 [119–136]
Trimester 1	42	109 [50–219]
Trimester 2	70	128 [88–219]
Trimester 3	64	172 [109–267]



**FIG. 4.** (A) Histogram of UI ( $\mu\text{g/L}$ ) for pregnant women in NHANES 2007–2010 ( $n=76$ ; reference line denotes the median UI =  $135.0 \mu\text{g/L}$ ). (B) Histogram of UI ( $\mu\text{g/L}$ ) for pregnant women ( $n=501$ ) in the NCS (reference line denotes the median UI =  $166.9 \mu\text{g/L}$ ). (C) Histogram of UI ( $\mu\text{g/L}$ ) for pregnant non-Hispanic black women, NHANES 2007–2010 ( $n=14$ ; reference line denotes the median UI =  $118.5 \mu\text{g/L}$ ). (D) Histogram of UI ( $\mu\text{g/L}$ ) for pregnant non-Hispanic black women ( $n=22$ ) in the NCS (reference line denotes the median UI =  $133 \mu\text{g/L}$ ).

contained iodine. Prenatal vitamins are not a dependable source of iodine, since only about 51% may contain iodine and of those that do, the measured iodine concentration can vary  $\pm 50\%$  from the labeled content (12). In addition to being an inconsistent source of iodine, relying on the use of prenatal vitamins to ensure iodine sufficiency during pregnancy may miss the very early stages of the fetal development. Not surprisingly, when we looked at supplement use (yes or no) in NHANES women of childbearing age, there was no significant correlation between those using supplements and those not using supplements.

There is a need to look beyond pregnancy status to identify what may be influencing UI concentrations in subpopulations. NHANES 2009–2010 indicates that non-Hispanic blacks had lower UI levels compared to non-Hispanic whites and all Hispanics across all age categories. However, the number of pregnant participants in NHANES was insufficient to look at pregnancy status by race/ethnicity. In the NCS pilot study, non-Hispanic black pregnant women ages 20 years and above had lower UI concentrations than both non-Hispanic whites and Hispanics, which was also lower than what WHO suggested for pregnant women. There were only five NCS

women less than 20 years, and their median UI was  $224 \mu\text{g/L}$  [CI 106–1701]. There is a wide confidence interval with a very small population.

The NHANES and NCS UI data suggest that non-Hispanic black women have lower UI concentration than other women. Additionally, non-Hispanic black women had lower dairy consumption, based on NHANES 2007–2008 data. Almost 40% of non-Hispanic black women of childbearing age never or only rarely consume dairy products compared with 27.1% of non-Hispanic white women and 31.1% of Hispanics. Moreover, proportionately more Hispanic and non-Hispanic white women of childbearing age reported consuming dairy products daily than do non-Hispanic blacks. Non-Hispanic black women reporting lower rates of dairy consumption is consistent with recent data on U.S. population reports of lactose intolerance (8). That study found that 76% of adults reporting to be lactose intolerant were female, and among females, 50% were non-Hispanic black, 30% non-Hispanic white, and 20% Hispanic. Self-diagnosed lactose intolerance and consequent avoidance of dairy products may be one of the contributing factors in the racial/ethnic differences we have shown in UI concentration.



TABLE 4. COMPARISON OF URINARY IODINE LEVELS BY RACE/ETHNICITY AND AGE GROUPS IN THE UNITED STATES

Age groups	Race/ethnicity	NHANES 2009–2010			NCS		
		n	UI ( $\mu\text{g/L}$ ) median [CI]	I/Cr ( $\mu\text{g/g creatinine}$ ) median [CI]	n	UI ( $\mu\text{g/L}$ ) median [CI]	I/Cr ( $\mu\text{g/g creatinine}$ ) median [CI]
6–11 years	Non-Hispanic white	109	241 [193–281]	285 [227–360]			
	Non-Hispanic black	84	168 [120–223]	163 [120–226]			
	All Hispanic	159	198 [153–254]	238 [216–268]			
12–19 years	Non-Hispanic white	164	141 [107–162]	125 [113–139]	— <sup>a</sup>		
	Non-Hispanic black	97	130 [107–155]	86 [72–109]	—		
	All Hispanic	162	150 [124–165]	132 [113–145]	—		
20+ years	Non-Hispanic white	951	143 [128–158]	147 [137–159]	327	179 [157–199]	208 [196–227]
	Non-Hispanic black	364	129 [117–144]	94 [84–101]	22	132 [104–221]	97 [86–189]
	All Hispanic	593	137 [125–156]	137 [126–144]	86	178 [136–209]	152 [135–177]
<i>Region</i>							
12+ years	North Carolina				82	217 [177–251]	169.4 [141.4–210.1]
	South Dakota/Minnesota				102	205 [161–225]	203.1 [177.9–242.7]
	Pennsylvania				45	125 [107–199]	196.4 [161.2–311.7]
	California				62	107 [86–147]	144.7 [121.4–195.2]
	New York				39	150 [86–186]	149.7 [126.3–189.6]
	Utah				102	190 [168–209]	197.5 [162.7–219.5]
	Wisconsin				69	145 [112–193]	241.2 [205.5–286.9]

<sup>a</sup>NCS 12–19 years has frequencies that fall below the disclosure threshold.

TABLE 5. URINARY IODINE MEDIAN AND 95% CONFIDENCE INTERVAL BY RACE AND DAIRY PRODUCT, SALT, AND FISH AND FISH/SHELLFISH CONSUMPTION IN WOMEN OF CHILDBEARING AGE, NHANES 2007–2008

Race	Dairy consumption	n	Weighted %	UI ( $\mu\text{g/L}$ ) median [CI]	p value <sup>a</sup>
Non-Hispanic white	Never or rare	149	27.1	111 [90–142]	0.0001
	Not often	167	31.1	131 [99–165]	
	Often	216	41.8	189 [163–204]	
Non-Hispanic black	Never or rare	126	39.8	111 [82–132]	0.0009
	Not often	89	27.9	109 [89–163]	
	Often	101	32.3	151 [110–212]	
All Hispanic	Never or rare	153	31.1	134 [106–155]	<0.0001
	Not often	133	27.2	163 [130–196]	
	Often	210	41.7	185 [155–207]	
<i>Salt consumption</i>					
Non-Hispanic white	Never or rare	142	27.1	110 [91–149]	0.07
	Not often	188	37.1	144 [116–189]	
	Often	183	35.9	148 [113–185]	
Non-Hispanic black	Never or rare	77	24.5	117 [93–161]	0.2
	Not often	97	29.5	132 [82–195]	
	Often	136	45.9	116 [97–132]	
All Hispanic	Never or rare	78	17.2	154 [124–239]	0.2
	Not often	128	27.1	152 [124–182]	
	Often	255	55.6	165 [141–194]	
<i>Fish/shellfish consumption</i>					
Non-Hispanic white	Yes	353	70.6	144 [114–178]	0.6
	No	159	29.4	134 [100–175]	
Non-Hispanic black	Yes	244	81.7	119 [99–151]	0.5
	No	66	18.3	123 [83–159]	
All Hispanic	Yes	339	73.4	166 [153–185]	0.03
	No	128	26.6	146 [108–185]	

<sup>a</sup>The *p* value tests whether the median UI concentrations were significantly different among never or rare, not often, and often dairy, and salt consumption groups within each race/ethnicity group. The *p* value tests whether the median UI concentrations were significantly different among yes or no answers for fish/shellfish consumption during the past 30 days within each race/ethnicity group.

Neither salt use during food preparation nor grain intake was significantly associated with an increase in median UI for any race/ethnic category of women of childbearing age. However, among women of childbearing age in NHANES who claimed to never or only rarely consume salt, Hispanic women had the highest median UI (154  $\mu\text{g}/\text{L}$ ) compared with non-Hispanic whites (110  $\mu\text{g}/\text{L}$ ) or non-Hispanic blacks (117  $\mu\text{g}/\text{L}$ ). The use of salt, consumption of grain, or fish/shellfish intake did not significantly affect UI, although there were racial/ethnic differences in reported salt use and fish/shellfish intakes.

Analysis has shown regional differences in UI among pregnant women in the NCS study. Future work with the NCS may provide the statistical power needed to stratify the participants within each region by race/ethnicity. The present data suggest that race/ethnicity is an important predictor of iodine status. However geographic location may prove to be an equally important predictor. Additionally, within pregnant women in the NCS, we have identified iodine insufficiency among non-Hispanic blacks and among NCS subgroups evaluated geographically. These data enhance ongoing efforts to identify populations at risk for iodine deficiency and may help direct efforts to ensure optimal fetal and postnatal development. NCS will be an important tool to assist in evaluation of U.S. population iodine intake in a vulnerable group. The NCS not only has important data on pregnant women but will collect data that may link the iodine status in pregnant women to birth and child health outcomes.

This study has several strengths. This is the largest known population of pregnant women to be surveyed for UI concentration at one time in the United States. Through NHANES and NCS we have a diverse sample set covering different ethnic groups, ages, geographic regions, and different stages of pregnancy. There are several limitations. These include the temporality of the UI biomarker, limited personal and health information, and unknown selection biases in the NCS convenience sample.

## Conclusion

The median UI concentration for the general U.S. population in NHANES 2009–2010 was significantly lower than the levels found during NHANES 2007–2008. These differences likely result from variations in dietary iodine intake for both the general population and for women of childbearing age. Grain, salt, and fish/shellfish had little effect on UI in women of childbearing age except for Hispanic women, in whom median UI was correlated with fish and shellfish intake. A higher proportion of non-Hispanic black women may avoid dairy products due to concerns about lactose intolerance. Children had a higher UI level than adults. Non-Hispanic white and Hispanic children have UI levels in the upper range of adequate or in the more than adequate range. Non-Hispanic black children have UI concentrations that put them solidly in the adequate range.

NCS data provide a unique look at a large population of pregnant women in the United States. The data indicate that third trimester pregnant non-Hispanic black women age 20 years and older were the only racial/ethnic group with median UI below 150  $\mu\text{g}/\text{L}$ . This racial/ethnic UI pattern was consistent with observations in NHANES data for the general population.

Understanding the median UI differences within subpopulations of the United States may help to identify where public health interventions should be focused to ensure adequate iodine nutrition for all, and can allow educational efforts to focus on groups for which evidence indicates inadequate iodine intake. Prevention of unnecessary self-imposed dairy restriction and improved iodine intake may be achieved by medical evaluation and education of individuals who perceive themselves to be lactose intolerant but who are not. Individuals with confirmed lactose intolerance, particularly women who are pregnant or who are of childbearing age, should be counseled about iodine supplementation. Public health intervention may include continued efforts by the medical community and public health officials to promote the use of iodine-containing prenatal vitamins and to promote continued efforts to monitor actual iodine content in prenatal vitamins. The NCS data have provided an opportunity to look across seven geographical locations, noting differences in median values among study sites. Further study is needed to determine if the concentration of iodine in the dairy consumed varies geographically and may be a significant factor in median UI concentrations, along with race/ethnicity.

## Acknowledgments

This analysis was conducted as part of the National Children's Study, supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), and funded, through its appropriation, by the Office of the Director of the National Institutes of Health. Supported by NICHD Contracts HHSN275200503414C, HHSN275200503411C, HHSN275200603416C, HHSN275200503415C, HHSN275200503413C, HHSN275200503410C, and HHSN275200503396C. The manuscript was developed by a Writing Team identified by the National Children's Study Publications Committee for the purpose of timely sharing of centrally collected NCS data. In addition, funding for the biospecimen analyses was provided by the National Center for Environmental Health, Centers for Disease Control and Prevention. The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention, the National Children's Study or the National Institutes of Health, or the U.S. Department of Health and Human Services.

## Disclosure Statement

The authors declare that no competing financial interests exist.

## References

1. Andersson M, Karumbunathan V, Zimmermann MB 2012 Global iodine status in 2011 and trends over the past decade. *J Nutr* **142**:744–750.
2. Hetzel B 1983 Iodine deficiency disorders (IDD) and their eradication. *Lancet* **322**:1126–1129.
3. Hetzel BS, Delange F, Dunn JT, Ling J, Mannar V, Pandav C 2004 Towards the global elimination of brain damage due to iodine deficiency. Oxford University Press, Delhi.
4. Bleichrodt N, Born M 1994 A meta analysis of research on iodine and its relationship to cognitive development. In: Stanbury J (ed) *The Damaged Brain of Iodine Deficiency*. Vol. Cognizant Communication, New York, pp 187–191.

5. Murray CW, Egan SK, Kim H, Beru N, Bolger PM 2008 US Food and Drug Administration's total diet study: dietary intake of perchlorate and iodine. *J Expo Sci Environ Epidemiol* **18**:571–580.
6. Hemken RW 1980 Milk and meat iodine content—relation to human health. *J Am Vet Med Assoc* **176**:1119–21.
7. Pearce EN, Pino S, He X, Bazrafshan HR, Lee SL, Braverman LE 2004 Sources of dietary iodine: bread, cows' milk, and infant formula in the Boston area. *J Clin Endocrinol Metab* **89**:3421–3424.
8. Nicklas TA, Qu HY, Hughes SO, He MY, Wagner SE, Foushee HR, Shewchuk RM 2011 Self-perceived lactose intolerance results in lower intakes of calcium and dairy foods and is associated with hypertension and diabetes in adults. *Am J Clin Nutr* **94**:191–198.
9. Dasgupta PK, Liu Y, Dyke JV 2008 Iodine nutrition: iodine content of iodized salt in the United States. *Environ Sci Technol* **42**:1315–1323.
10. Institute of Medicine 2001 Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. The National Academies Press, Washington, DC.
11. Public Health Committee of the American Thyroid Association, Becker DV, Braverman LE, Delange F, Dunn JT, Franklyn JA, Hollowell JG, Lamm SH, Mitchell ML, Pearce E, Robbins J, Rovet JF 2006 Iodine supplementation for pregnancy and lactation—United States and Canada: recommendations of the American Thyroid Association. *Thyroid* **16**:949–951.
12. Gregory CO, Serdula MK, Sullivan KM 2009 Use of supplements with and without iodine in women of childbearing age in the united states. *Thyroid* **19**:1019–1020.
13. World Health Organization 2008 Assessment of iodine deficiency disorders and monitoring their elimination—a guide for programme managers. Third edition. World Health Organization, Geneva.
14. Caldwell KL, Makhmudov A, Ely E, Jones RL, Wang RY 2011 Iodine status of the U.S. population, National Health and Nutrition Examination Survey, 2005–2006 and 2007–2008. *Thyroid* **21**:419–427.
15. Mortensen M, Hirschfeld S 2012 The National Children's Study: an opportunity for medical toxicology. *J Med Toxicol* **8**:160–165.
16. The National Children's Study. What is the National Children's Study? Available at [www.nationalchildrensstudy.gov/Pages/default.aspx](http://www.nationalchildrensstudy.gov/Pages/default.aspx) (accessed August 3, 2012).
17. Caldwell KL, Jones R, Hollowell JG 2005 Urinary iodine concentration: United States National Health and Nutrition Examination Survey 2001–2002. *Thyroid* **15**:692–699.
18. Caldwell KL, Maxwell CB, Makhmudov A, Pino S, Braverman LE, Jones RL, Hollowell JG 2003 Use of inductively coupled plasma mass spectrometry to measure urinary iodine in NHANES 2000: comparison with previous method. *Clin Chem* **49**:1019–1021.
19. Caudill SP, Schleicher RL, Pirkle JL 2008 Multi-rule quality control for the age-related eye disease study. *Stat Med* **27**:4094–4106.
20. Lin LI 1989 A concordance correlation coefficient to evaluate reproducibility. *Biometrics* **45**:255–268.
21. Centers for Disease Control and Prevention. Ensuring the quality of urinary iodine procedures. Available at [www.cdc.gov/labstandards/equip.html](http://www.cdc.gov/labstandards/equip.html) (accessed August 3, 2012).
22. Korn EL, Graubard BI 1998 Confidence intervals for proportions with small expected number of positive counts estimated from survey data. *Surv Methodol* **24**:193–201.
23. Williams RL, Perritt RL 1986 The Weighted Median Test for Finite Population Samples. American Statistical Association, Washington, DC.

Address correspondence to:

*Kathleen L. Caldwell, PhD*

*Division of Laboratory Sciences*

*National Center for Environmental Health*

*Centers for Disease Control and Prevention*

*4770 Buford Highway, NE, Mail Stop F-18*

*Atlanta, GA 30341*

*E-mail: klc7@cdc.gov*